

Chicago Flu Update CHICAGO Immunization Program

April 29, 2016

Chicago Department of Public Health Julie Morita, MD, Commissioner

What is the risk?

Rahm Emanuel, Mayor

Currently, the risk of influenza infection continues to decrease. Vaccination is the best way to protect against influenza infection and all Chicagoans aged six months and older are encouraged to get vaccinated. Chicagoans should ask their healthcare provider or pharmacist about vaccine availability. For those without a healthcare provider or whose healthcare providers do not have the influenza vaccine, a list of <u>City of Chicago Fast-Track Immunization</u> <u>Clinics</u>¹ is available on the City website and by calling 311. To locate the closest City of Chicago clinic or retail pharmacy, go to <u>www.chicagoflushots.org</u>.

Are severe cases of influenza occurring?

The Illinois Department of Public Health (IDPH) has issued <u>influenza testing and reporting recommendations</u>². Suspected novel and variant influenza, pediatric influenza-associated deaths, influenza-associated ICU hospitalizations and outbreaks of influenza-like illness in a congregate setting should all be reported to CDPH via IN-EDSS³. For the week of April 17-23, 2016 (week 16), 5 influenza-associated ICU hospitalizations were reported (**Figure 1**).

Since October 4, 2015, 264 influenza-associated ICU hospitalizations have been reported; 249 were positive for influenza A (8 H3N2, 138 (H1N1)pdm09 and 103 unknown subtype [subtyping not attempted or not all subtypes tested]) and 15 were positive for influenza B. One hundred twenty-six (48%) were non-Hispanic Black, 151 (57%) were male, and 155 (59%) were 50 years of age or older (median age of 54 years with a range of 3 months-99 years). Seventy-eight (30%) had reported lung disease (e.g. asthma, COPD), 68 (26%) had active cardiac disease and 75 (28%) required ventilator support. Seventeen deaths were reported among ICU admissions. Five outbreaks consisting of two or more confirmed cases of influenza were reported in long-term care facilities.

How much influenza-like illness is occurring? CDPH receives data from influenza surveillance sites across Chicago, which report the total number of patient visits seen weekly, and of those visits, the number with influenza-like illness (ILI) (i.e., fever of 100°F or greater, with cough or sore throat). All hospitals in Chicago that provide emergent care are required to report on a weekly basis the total number of emergency department visits, and of those visits, the number with ILI. For the week of April 17-23, 2016 with 10 hospitals reporting, **3.6%** of emergency department visits were due to ILI (**Figure 2**).

Figure 1. Number of influenza-associated intensive care unit hospitalizations reported for Chicago residents, for current season (2015-2016) and previous season (2014-2015), October-May.



Figure 2. Percent of <u>emergency department</u> visits attributed to influenza-like illness based on manual reporting as determined by individual hospitals, Chicago, by week, for the current season (2015-2016) and previous three seasons, October-May.



ESSENCE is an electronic syndromic surveillance system that utilizes emergency department chief complaint data submitted daily by participating Chicago hospitals. ILI activity is determined solely based on the patient's chief complaint and does not take into account the entire medical record, as the ILI activity reported in Figure 2 does. Currently, 10 Chicago hospitals submit data to ESSENCE. For the week of April 17-23, 2016, **1.3%** of emergency department visits were due to ILI (**Figure 3**).

in CDC's Influenza-like Illness Surveillance Network (ILINet) by reporting on the number of patients with ILI seen weekly. From April 17-23, 2016, with 24 facilities reporting, 3.3% of visits were due to influenza-like illness (Figure 4).

Which influenza strains are circulating?

Data on influenza virus test results are reported by Chicago laboratories performing influenza subtyping. For the week of April 17-23, 2016, with 5 laboratories reporting, 31 of the 432 (7%) specimens tested for influenza were positive (14 A (H1N1)pdm09, 1 A (H3N2), 3 A [unknown subtype] and 13 influenza B). Since October 4, 2015, 1,500 of 18,308 (8%) specimens tested for influenza have been positive; 1,388 typed as influenza A (40 H3N2, 1,111 (H1N1)pdm09, 237 unknown subtype [subtyping not attempted or not all subtypes tested]) and 112 were typed as influenza B (Figure 5). The cumulative percent of specimens testing positive for influenza is lower than last season during the same time period (Table 1).

Table 1. Cumulative percent of specimens testing positive for influenza by subtype for the current season (2015-2016) and the previous five seasons, Chicago, Weeks 40-16.

Influenza Season	Total Positive	A H3N2	A H1N1pdm09	A Not Subtyped	В
2010-2011	11%	26%	24%	27%	23%
2011-2012	8%	15%	17%	16%	51%
2012-2013	13%	66%	6%	13%	15%
2013-2014	8%	3%	62%	22%	13%
2014-2015	14%	69%	0%	13%	28%
2015-2016	8%	3%	74%	16%	7%

Where can I get more information?

The Centers for Disease Control and Prevention's FluView⁴ report provides national updates and trends reed States, as well as detailed information on antiviral resistance, severity of illness, and other topics. Updates specific to <u>Illinois</u>⁵ and <u>Suburban Cook County</u>⁶ are also available online. Current and archived issues of the Chicago Flu Update can be found on the CDPH website section <u>Current Flu Situation in Chicago</u>⁷. In 2013, the Metropolitan Chicago Healthcare Council (MCHC) and CDPH released "Stop the Spread: A Health Care Guide to Influenza Preparedness³⁸. This report provides an overview of influenza, its impact on public health and how hospitals can prepare for, mitigate the impact of and respond to influenza infections and outbreaks.

> The 21st Annual Chicago Infection Control Conference May 20th, 2016 at the Chicago Cultural Center 78 E. Washington St. Chicago, 60602 **Registration is Now Open!**

Several outpatient clinics throughout Chicago participate Figure 3. Percent of emergency department visits attributed to influenza-like illness based on chief complaint data submitted to ESSENCE, Chicago, by week, for the current season (2015-2016) and the previous three seasons, October-May.



Figure 4. Percent of medically attended outpatient visits attributed to influenza-like illness as reported by ILINet facilities, Chicago, by week, for the current season (2015-2016) and the previous three seasons, October-May.



lated to the intensity of influenza activity across the Unit- Figure 5. Percent of specimens testing positive (by RT-PCR) for influenza by subtype as reported by local laboratories serving Chicago hospitals, for the current season (2015-2016) October-May.



All data are preliminary and may change as more reports are received

4http://www.cdc.gov/flu/weekly/index.htm; 5http://www.idph.state.il.us/flu/surveillance.htm; 6http://www.cookcountypublichealth.org/data-reports#Influenza; 7http:// www.cityofchicago.org/city/en/depts/cdph/supp_info/influenza/current_flu_situationinchicago2011.html; 8 https://www.mchc.com/wp-content/uploads/2014/11/ MCHC Influenza 2013.pdf