

Chicago Flu Update



City of Chicago Rahm Emanuel, Mayor

November 28, 2014

Chicago Department of Public Health Bechara Choucair, MD, Commissioner

What is the risk?

Currently, the risk of influenza infection has increased slightly. Vaccination is the best way to protect against influenza infection and all Chicagoans aged six months and older are encouraged to get vaccinated. Chicagoans should ask their healthcare provider or pharmacist about vaccine availability. Vaccinate Illinois Week is a localized observance of National Influenza Vaccination Week which is scheduled for December 7th through the 13th. CDPH and partners throughout the state will be reminding providers and the public that it is not too late to give or get an influenza vaccine. For those without a healthcare provider or whose healthcare providers do not have the influenza vaccine, a schedule of City of Chicago influenza vaccine clinics is available on the City website and by calling 311. To locate the closest City of Chicago clinic or retail pharmacy, go to www.chicagoflushots.org.

Are severe cases of influenza occurring?

The Illinois Department of Public Health (IDPH) has issued <u>influenza testing and reporting guidelines</u>². Suspected novel and variant influenza, pediatric influenza-associated deaths, influenza-associated ICU hospitalizations and outbreaks of influenza-like illness in a congregate setting should all be reported to CDPH via <u>IN-EDSS</u>³. For the week of November 16-22, 2014, one influenza-associated ICU hospitalization was reported which was positive for influenza A (H3N2).

Since September 28, 2014, five influenza-associated ICU hospitalizations have been reported (**Figure 1**). Among the total ICU hospitalizations reported 4 were positive for influenza A (3 H3N2 and 1 unknown subtype) and 1 was positive for influenza B.

How much influenza-like illness is occurring?

CDPH receives data from over 60 surveillance sites across Chicago, which report the total number of patient visits seen weekly, and of those visits, the number with influenza-like illness (ILI) (i.e., fever of 100°F or greater, with cough or sore throat). All hospitals in Chicago that provide emergent care are required to report on a weekly basis the total number of emergency department visits, and of those visits, the number with ILI. For the week of November 16-22, 2014 (week 47), with 13 hospitals reporting, 3.5% of emergency department visits were due to ILI (**Figure 2**).

ESSENCE is an electronic syndromic surveillance system that utilizes emergency department chief complaint data submitted daily by participating Chicago hospitals. ILI activity is determined solely based on the patient's chief complaint and does not take into account the entire medical record, as the ILI activity reported in Figure 2 does.

Currently, 10 Chicago hospitals submit data to ES-

SENCE. For the week of November 16-22, 2014, 1.8% of emergency department visits were due to ILI (**Figure 3**). Several outpatient clinics and hospital emergency departments throughout Chicago participate in CDC's Influenza-like Illness Surveillance Network (**ILINet**) by reporting on the number of patients with ILI seen weekly. From November 9-15, 2014, with 42 facilities reporting, 3.4% of visits were due to influenza-like illness (**Figure 4**).

Figure 1. Number of influenza-associated intensive care unit hospitalizations reported for Chicago residents, for current season (2014-2015) and previous season (2013-2014), October-May.





Figure 2. Percent of <u>emergency department</u> visits attributed to influenza-like illness based on manual reporting as determined by individual hospitals, Chicago, by week, for the current season (2014-2015) and previous three seasons, October-May.



Which influenza strains are circulating?

Data on influenza virus test results are reported by Chicago laboratories performing influenza subtyping. For the week of November 16-22, 2014, with 6 laboratories reporting, 32 of the 379 (8.4%) specimens tested for influenza were positive. During the same week last season, 2.4% of specimens tested were positive for influenza. Among this week's positive specimens, 29 were typed as influenza A (25 H3N2 and 4 unknown subtype) and 3 was typed as influenza B. Since September 28, 2014, 57 of 2,781 (2%) specimens tested for influenza have been positive; 45 typed as influenza A (38 H3N2 and 7 unknown subtype) and 12 were typed as influenza B. (Figure 5).

Influenza A (H3N2) Antigenic Drift

Since October 1, 2014, CDC has characterized 85 influenza A (H3N2) viruses. Of these, 41 (48%) are like the A (H3N2) component of the Northern Hemisphere influenza vaccine. Forty-four (52%) of the 85 viruses tested have been characterized as an antigenic variant ("drifted") virus which has been selected for the 2015 Southern Hemisphere influenza vaccine. If drifted influenza A (H3N2) viruses circulate broadly in the U.S. this season, this could translate into a reduced vaccine effectiveness against circulating H3N2 viruses. However, studies have shown evidence that seasonal influenza vaccination can sometimes induce cross-protection. While the vaccine effectiveness may be reduced, the vaccine can still offer protection. In addition, more than one type of influenza usually circulates during a single season and influenza vaccines protect against three or four different influenza viruses. Vaccination continues to offer the best protection against influenza infection even when there are some antigenically drifted viruses circulating in the community.

Where can I get more information?

The Centers for Disease Control and Prevention's FluView⁴ report provides national updates and trends related to the intensity of influenza activity across the United States, as well as detailed information on antiviral resistance, severity of illness, and other topics. Updates specific to Illinois⁵ and Suburban Cook County⁶ are also available online. Current and archived issues of the *Chicago Flu Update* can be found on the CDPH website section Current Flu Situation in Chicago⁷. In 2013, the Metropolitan Chicago Healthcare Council (MCHC) and CDPH released "Stop the Spread: A Health Care Guide to Influenza Preparedness". This report provides an overview of influenza, it's impact on public health and how hospitals can prepare for, mitigate the impact of and respond to influenza infections and outbreaks.

Figure 3. Percent of <u>emergency department</u> visits attributed to influenza-like illness based on chief complaint data submitted to <u>ESSENCE</u>, Chicago, by week, for the current season (2014-2015) and the previous three seasons, October-May.




Figure 4. Percent of medically attended visits attributed to influenza -like illness as reported by ILINet facilities, Chicago, by week, for the current season (2014-2015) and the previous three seasons, October-May.




Figure 5. Percent of specimens testing positive (by RT-PCR) for influenza by subtype as reported by local laboratories serving Chicago hospitals, for the current season (2014-2015) October-May,

