

Pfizer-BioNTECH **COVID-19 Vaccine**

Ages: 16 years of age and older

Use for: Any dose in the 2-dose series.

> COVID-19 vaccines are NOT interchangeable. Both doses MUST be COVID-19 vaccine (Pfizer).

Route: Intramuscular (IM) injection

> Prior to administration, mix with 0.9% sodium chloride (normal saline, preservative-free)

diluent ONLY.

Beyond

Use Time: Use within 6 hours

of mixing.



Pfizer-BioNTECH **COVID-19 Vaccine**

Ages: 16 years of age and older

Use for: Any dose in the 2-dose series.

COVID-19 vaccines are NOT interchangeable. Both doses MUST be COVID-19 vaccine (Pfizer).

Route: Intramuscular (IM) injection

> Prior to administration, mix with 0.9% sodium chloride (normal saline, preservative-free)

diluent ONLY.

Beyond

Use Time: Use within 6 hours

of mixing.





Pfizer-BioNTECH **COVID-19 Vaccine**

Ages: 16 years of age and older

Use for: Any dose in the 2-dose series.

> COVID-19 vaccines are NOT interchangeable. Both doses MUST be COVID-19 vaccine (Pfizer).

Route: Intramuscular (IM) injection

> Prior to administration, mix with 0.9% sodium chloride (normal saline, preservative-free)

diluent ONLY.

Bevond

Use Time: Use within 6 hours

of mixing.



Pfizer-BioNTECH COVID-19 Vaccine

Ages: 16 years of age and older

Use for: Any dose in the 2-dose series.

> COVID-19 vaccines are NOT interchangeable. Both doses MUST be COVID-19 vaccine (Pfizer).

Route: Intramuscular (IM) injection

> Prior to administration, mix with 0.9% sodium chloride (normal saline, preservative-free)

diluent ONLY.

Beyond

Use Time: Use within 6 hours

of mixing.



Pfizer-BioNTECH **COVID-19 Vaccine**

16 years of age and older Ages:

Use for: Any dose in the 2-dose series.

COVID-19 vaccines are NOT interchangeable. Both doses MUST be COVID-19 vaccine (Pfizer).

Route: Intramuscular (IM) injection

> Prior to administration, mix with 0.9% sodium chloride (normal saline, preservative-free)

diluent ONLY.

Beyond

Use Time: Use within 6 hours

of mixing.



Pfizer-BioNTECH **COVID-19 Vaccine**

Ages: 16 years of age and older

Use for: Any dose in the 2-dose series. COVID-19 vaccines are NOT interchangeable.

Both doses MUST be COVID-19 vaccine (Pfizer).

Route: Intramuscular (IM) injection

> Prior to administration, mix with 0.9% sodium chloride (normal saline, preservative-free)

diluent ONLY.

Beyond

Use Time: Use within 6 hours

of mixing.



12/12/20